

---

## Ju The Bmj

**use of quick response (qr) coded ... - innovationsj** - sellick jfk, weaver ju. *bmj innov* 2018;4:115-122. health it, systems and process innovations abstract background an adrenal crisis is a rare but life-threatening condition in which hydrocortisone and fluid resuscitation must be given promptly to prevent hypotensive shock and death. a clinical audit assessed healthcare professionals' **cardiovascular safety of methylphenidate among ... - bmj** - ju-young shin, 1, 2 elizabeth e roughead, 3 byung-joo park, 4 nicole l pratt abstract objective to determine whether treatment with methylphenidate in children and young people with attention-deficit/hyperactivity disorder (adhd) was associated with ... *bmj* 2016;353:i2550 | doi: 10.1136/bmj.i2550 **analysis research methodology reasons or excuses ... - bmj** - *bmj* | 21 ne 2008 ju | volu 336 me 1413 analysis heterogeneous data are a common problem in meta-analysis. john ioannidis, nikolaos patsopoulos, and . hannah rothstein. show that final synthesis is possible and desirable in most cases . reasons or excuses for avoiding meta-analysis in forest plots. some systematic reviews simply assemble **bmj ebm: first published as 10.1136/bmjebm-2018-111012 on ... - bmj ebm: first published as 10.1136/bmjebm-2018-111012 on 27 ju ... bmj evidence-based medicine article criticizing the cochrane review of hpv vaccines on 27th july 2018, an article was published in the journal bmj evidence-based medicine relating to the recently published cochrane review on prophylactic human papillomavirus (hpv) vaccines ... the future of the medical profession - jcpj - bmj: first published as 10.1136/bmj.39265.679375.80 on 12 july 2007. downloaded from . editorials. 54. bmj | 14 ly 2007 ju | volu 335me. diarrhoea associated with antibiotic use. evidence supports the use of probiotics, but effectiveness depends on the strain. diarrhoea is a common side effect of antibiotics; it . **research methods of the international tobacco control (itc ... - were not surveyed by waves 1 and 2; two ju wei hui were selected within the new jie dao, with probability proportional to ju wei hui population size. 2. if only one ju wei hui was needed at wave 3 from the new jie dao, the jie dao was first divided in half in terms of population (depending on the number of ju wei hui in the jie dao). fine mapping mhc associations in graves ... - jmgj** - by bmj. abstract background the classical human leucocyte antigen (hla) genes were the most important genetic determinant for graves' disease (gd). the aim of the study was to fine map causal variants of the hla genes. methods we applied imputation with a pan-asian hla reference panel to thoroughly investigate the major **bdc e fhgjjkjlNm\*g - staff.washington** - d! fa ^czy[aj ztx9bc`izy[\z]r\_`a`Sitemap | [Best Seller](#) | [Home](#) | [Random](#) | [Popular](#) | [Top](#)**